




How to administer PREVACID SoluTab by nasogastric (NG) tube or oral syringe

Administration of PREVACID SoluTab by NG tube (for patients who have difficulty swallowing tablets and have a ≥ 8 French NG tube in place)¹



1. Place PREVACID SoluTab in syringe.



3. Shake gently to allow for quick dispersal.



2. Draw up water into syringe.
– For 15 mg tablet, draw up 4 mL of water.
– For 30 mg tablet, draw up 10 mL of water.



4. After the tablet has dispersed, inject through the NG tube into the stomach within 15 minutes.

5. Refill the syringe with approximately 5 mL of water, shake gently, and flush the NG tube.

By oral syringe (for patients who have difficulty swallowing tablets)¹



1. Place PREVACID SoluTab in oral syringe.



3. Shake gently to allow for quick dispersal.



2. Draw up water into syringe.
– For 15 mg tablet, draw up approximately 4 mL of water.
– For 30 mg tablet, draw up approximately 10 mL of water.



4. After tablet has dispersed, administer contents orally within 15 minutes.

5. Refill the syringe with approximately 2 mL of water for the 15 mg tablet or 5 mL for the 30 mg tablet, shake gently, and administer any remaining contents.

PREVACID products should not be crushed or chewed.

Phenylketonurics: contains phenylalanine 2.5 mg per 15 mg tablet and 5.1 mg per 30 mg tablet.

For further information, including adverse events, please see the reverse side.